

# QUATERNIONIC ANALYSIS

SUNAY MIDUTHURI

**ABSTRACT.** In this paper, we discuss the topic of Quaternionic Analysis, in which we attempt to extend complex analysis to the quaternions.

## 1. INTRODUCTION

Quaternions are a four-dimensional algebra that extend the idea of a complex numbers to create an entirely new number system. They were invented by the Irish mathematician William Rowan Hamilton in [Ham66]. Much of the modern theory of Quaternionic Analysis was developed by Fueter in [Fue35].

**Definition 1.1.** The set of quaternions, denoted  $\mathbb{H}$ , is a 4-dimensional vector space over  $\mathbb{R}$  with basis  $1, i, j, k$ , where  $i, j, k$  are the elementary quaternions.

**Definition 1.2.** The elementary quaternions satisfy

$$i^2 = j^2 = k^2 = ijk = -1.$$

Furthermore, they are anticommutative, meaning that

$$ij = -ji.$$

From this, we trivially obtain the result of Theorem 1.3.

**Theorem 1.3.** *The elementary quaternions satisfy*

$$ij = k, jk = i, ki = j.$$

We are now ready to extend the elementary quaternions to the larger set of quaternions.

**Definition 1.4.** A quaternion can be represented as a linear combination in the form

$$q = w + xi + yj + zk,$$

where  $x, y, z \in \mathbb{R}$ .

**Definition 1.5.** The conjugate of a quaternion  $q = w + xi + yj + zk$  is

$$\bar{q} = w - xi - yj - zk.$$

**Definition 1.6.** The magnitude of a quaternion  $q = w + xi + yj + zk$  is

$$|q| = \sqrt{w^2 + x^2 + y^2 + z^2}.$$

**Definition 1.7.** If  $q = t + xi + yj + zk \in H$ , then the real part  $\text{Re}(q) = t$  and the pure quaternion part  $\text{Pu}(q) = xi + yj + zk$ .

The quaternions  $\mathbb{H}$  form a 4-dimensional algebra over  $\mathbb{R}$ . If  $q$  is any quaternion linearly independent from 1, the vector subspace spanned by  $q$  and 1 is a subfield of  $\mathbb{H}$  that is isomorphic to  $\mathbb{C}$  [Sud98]. We can also embed  $\mathbb{C}$  in  $\mathbb{H}$ ; for instance, any quaternion  $q \in \mathbb{H}$  can be expressed in the form

$$q = v + jw,$$

where  $v, w \in \mathbb{C}$  and  $j$  is an elementary quaternion.

## 2. DIFFERENTIAL FORMS

**Definition 2.1.** We define the differential of a function  $f : \mathbb{H} \rightarrow \mathbb{H}$  as

$$df = \frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy + \frac{\partial f}{\partial z} dz.$$

If this exists, we say that  $f$  is *real-differentiable*.

**Definition 2.2.** The differential of  $f : \mathbb{H} \rightarrow \mathbb{H}$  at a point  $q$  is then defined analogously to Definition 2.1 as a linear mapping  $df_q : \mathbb{H} \rightarrow \mathbb{H}$ .

**Definition 2.3.** We also define the quaternion gradient operator as

$$\square = \frac{\partial}{\partial w} + i \frac{\partial}{\partial x} + j \frac{\partial}{\partial y} + k \frac{\partial}{\partial z}.$$

**Definition 2.4.** Finally, we define the differential of the identity function

$$dq = dt + i dx + j dy + k dz.$$

## 3. FUNCTIONS

**Definition 3.1.** A quaternion function  $f : \mathbb{H} \rightarrow \mathbb{H}$  is quaternion-differentiable on the left at a point  $q$  if

$$\frac{df}{dq} = \lim_{h \rightarrow 0} \frac{f(q+h) - f(q)}{h}$$

exists as  $h \rightarrow 0$  from any direction in  $\mathbb{H}$ .

**Theorem 3.2.** Suppose that a function  $f : \mathbb{H} \rightarrow \mathbb{H}$  is defined and quaternion-differentiable on the left throughout a connected open set  $U$ . Then on  $U$ ,  $f$  has the form

$$f(q) = aq + b,$$

where  $a, b \in \mathbb{H}$ .

*Proof.* [Sud98] If  $f$  is quaternion-differentiable on the left, we can write

$$df_q(h) = h \frac{df}{dq} = dq \frac{df}{dq}.$$

We then equate coefficients of  $t, x, y, z$  to obtain

$$\begin{aligned}\frac{df}{dq} &= \frac{\partial f}{\partial t} \\ &= -i \frac{\partial f}{\partial x} \\ &= -j \frac{\partial f}{\partial y} \\ &= -k \frac{\partial f}{\partial z}.\end{aligned}$$

Now suppose that  $q = v + jw$ , where  $v = t + ix$  and  $w = y - iz$ , and let  $f(q) = g(v, w) + jh(v, w)$ , where  $g$  and  $h$  are complex functions of  $v$  and  $w$ . We can then separate our the previous equation into two sets of equations:

$$\frac{\partial g}{\partial t} = -i \frac{\partial g}{\partial x} = \frac{\partial h}{\partial y} = i \frac{\partial h}{\partial z},$$

and

$$\frac{\partial h}{\partial t} = i \frac{\partial h}{\partial x} = -\frac{\partial g}{\partial y} = i \frac{\partial g}{\partial z}.$$

We can write these in terms of complex derivatives as

$$\begin{aligned}\frac{\partial g}{\partial \bar{v}} &= \frac{\partial h}{\partial \bar{w}} = \frac{\partial h}{\partial v} = \frac{\partial g}{\partial w} = 0, \\ \frac{\partial g}{\partial v} &= \frac{\partial h}{\partial w},\end{aligned}$$

and

$$\frac{\partial h}{\partial \bar{v}} = -\frac{\partial g}{\partial \bar{w}}.$$

Then, by Cauchy-Riemann,  $g$  is analytic in  $v$  and  $\bar{w}$ , and  $h$  is analytic in  $\bar{v}$  and  $w$ . Then

$$\frac{\partial^2 g}{\partial v^2} = \frac{\partial g}{\partial v} \left( \frac{\partial h}{\partial w} \right) = \frac{\partial g}{\partial w} \left( \frac{\partial h}{\partial v} \right) = 0,$$

from which we find that  $g$  is linear and the desired result follows.  $\square$

**Definition 3.3.** Conformal mappings in quaternionic analysis, as in complex analysis, are those which preserve angles.

#### 4. REGULAR FUNCTIONS

**Definition 4.1.** We define the Left Cauchy-Riemann-Fueter equation of a function  $f : \mathbb{H} \rightarrow \mathbb{H}$  as

$$\frac{\partial_l f}{\partial \bar{q}} = \frac{\partial f}{\partial t} + i \frac{\partial f}{\partial x} + j \frac{\partial f}{\partial y} + k \frac{\partial f}{\partial z}.$$

**Definition 4.2.** Similarly, we define the Right Cauchy-Riemann-Fueter equation of a function  $f : \mathbb{H} \rightarrow \mathbb{H}$  as

$$\frac{\partial_r f}{\partial \bar{q}} = \frac{\partial f}{\partial t} + \frac{\partial f}{\partial x} i + \frac{\partial f}{\partial y} j + \frac{\partial f}{\partial z} k.$$

**Definition 4.3.** A function  $f : \mathbb{H} \rightarrow \mathbb{H}$  is left-regular or right regular if

$$\frac{\partial_l f}{\partial \bar{q}} = 0$$

or

$$\frac{\partial_r f}{\partial \bar{q}} = 0,$$

respectively.

If we write  $q = v + jw$ , the equations simplify to

$$\frac{\partial g}{\partial \bar{v}} = \frac{\partial h}{\partial \bar{w}}, \quad \frac{\partial g}{\partial w} = -\frac{\partial h}{\partial v}.$$

**Definition 4.4.** A harmonic function  $f : \mathbb{H} \rightarrow \mathbb{H}$  is one that satisfies

$$\frac{\partial^2 f}{\partial t^2} + \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} = 0.$$

**Definition 4.5.** We define

$$\Gamma_r(df) = \frac{\partial f}{\partial t} + i \frac{\partial f}{\partial x} + j \frac{\partial f}{\partial y} + k \frac{\partial f}{\partial z}.$$

**Theorem 4.6.** *The left differential  $\partial_l$  can be written as*

$$\partial_l(f) = \frac{1}{2} \bar{\Gamma}_r(df).$$

**Theorem 4.7.** *[Sud98] If  $f$  is a harmonic function, then*

$$\partial_l f$$

*is regular.*

**Theorem 4.8.** *[Sud98] Let  $u$  be a real-valued function defined on a star-shaped open set  $U \subseteq \mathbb{H}$ . If  $U$  is harmonic and has a continuous second derivative, there exists a regular function  $f$  defined on  $U$  such that  $\text{Re}(f) = u$ .*

## 5. INTEGRALS

**Definition 5.1.** We define the 3-form

$$Dq(x, y, z)$$

to be the alternating trilinear form

$$\langle w, Dq(x, y, z) \rangle = v(w, x, y, z),$$

where the volume form

$$v = dt \wedge dx \wedge dy \wedge dz.$$

Geometrically,  $Dq(x, y, z)$  is the quaternion perpendicular to  $x, y, z$  with magnitude equal to the volume of the parallelepiped given by  $x, y, z$ .

**Theorem 5.2.** *[Sud98]  $Dq(a, b, c) = \frac{1}{2}(c\bar{a}b - b\bar{a}c)$ .*

**Theorem 5.3.** *[Sud98] A differentiable function  $f$  is regular at  $q$  if and only if*

$$Dq \wedge df_q = 0.$$

From this result, we find that

$$\int_{\partial D} Dq f = 0$$

if  $f$  is regular and continuously differentiable on a set  $D$ .

We also obtain the Cauchy-Fueter Integral Formula:

**Theorem 5.4.** *If  $f$  is regular on every point of a 4-parallelopiped  $C$  and  $q_0 \in C$ ,*

$$f(q_0) = \frac{1}{2\pi^2} \int_{\partial C} \frac{(q - q_0)^{-1}}{|q - q_0|^2} Dq f(q).$$

Finally, we get to Cauchy's theorem:

**Theorem 5.5.** *Suppose  $f$  is regular in an open set  $U$ , and let  $C$  be a rectifiable 3-chain which is homologous to 0 in the singular homology of  $U$ . Then*

$$\int_C Dq f = 0.$$

We end with one application of quaternions to number theory.

## 6. LAGRANGE'S FOUR-SQUARE THEOREM

In this section, we examine Lagrange's Four-Squares Theorem and its proof using quaternionic analysis. The theorem statement is as follows:

**Conjecture 6.1** (Lagrange's Four-Square Theorem). *Every nonnegative integer  $p$  can be written as*

$$p = a^2 + b^2 + c^2 + d^2,$$

*where  $a, b, c$ , and  $d$  are nonnegative integers.*

To prove this, we'll first need several definitions to develop the necessary theory.

**Definition 6.2.** A Lipschitz quaternion is a quaternion of the form

$$a + bi + cj + dk,$$

where  $a, b, c, d \in \mathbb{Z}$ .

**Definition 6.3.** The Hurwitz Integers are the set of all quaternions with all-integer or all-half-integer components; that is, any Hurwitz Integer  $h$  can be expressed in the form

$$h = \frac{1}{2}a(1 + i + j + k) + bi + cj + dk,$$

where  $a, b, c$  are all integers.

The following theorem then follows trivially.

**Theorem 6.4.** *The sum of any two Hurwitz Integers is itself a Hurwitz integer.*

**Definition 6.5.** A Hurwitz integer  $a$  is a multiple of a Hurwitz integer  $b$  if and only if there is a Hurwitz integer  $c$  such that  $a = bc$ .

**Definition 6.6.** [CW11] A Hurwitz Prime is a Hurwitz Integer which has no non-trivial factors (that is, factors other than elementary quaternions and

$$\pm \frac{1}{2} \pm \frac{1}{2}i \pm \frac{1}{2}j \pm \frac{1}{2}k$$

).

From now on, we'll refer to typical primes  $p \in \mathbb{Z}$  as "regular primes", and Hurwitz Primes as "Hurwitz Primes" to clearly differentiate the two.

**Lemma 6.7.** [CW11] *If positive integers  $p$  and  $q$  can be written as the sum of four squares, so can their product.*

*Proof.* Suppose that  $p = a^2 + b^2 + c^2 + d^2$  and  $q = w^2 + x^2 + y^2 + z^2$ . Then, if

$$\alpha = a + bi + cj + dk \text{ and } \beta = w + xi + yj + zk,$$

we know that  $p = \|\alpha\|^2$  and  $q = \|\beta\|^2$ , for some Lipschitz quaternions  $\alpha$  and  $\beta$ . Then  $uv = \|\alpha\|^2 \|\beta\|^2 = \|\alpha\beta\|^2 = \|A\|^2$  for some Lipschitz quaternion  $A$ , from which the desired result follows.  $\square$

This means that it suffices to show the Lagrange Four-Square Theorem for all regular primes. The following two lemmas from [CW11] are given without proof, as said proofs are not particularly relevant to the Lagrange Four-Square Theorem.

**Lemma 6.8.** *Let  $p$  be a Hurwitz prime. If*

$$p \mid \alpha\beta,$$

*either  $p \mid \alpha$  or  $p \mid \beta$ .*

**Lemma 6.9.** *If  $p$  is an odd regular prime, there exist integers  $l$  and  $m$  such that*

$$p \mid 1 + l^2 + m^2.$$

**Theorem 6.10.** *If  $p$  is an odd regular prime, it is not a Hurwitz Prime.*

*Proof.* We'll use contradiction. Suppose that  $p$  is a Hurwitz prime. Now, by Lemma 6.9, there exist integers  $l$  and  $m$  such that

$$p \mid 1 + l^2 + m^2.$$

But

$$\begin{aligned} (1 + li + mj)(1 - li - mj) &= (1 + (li + mj))(1 - (li + mj)) \\ &= (1)^2 - (li + mj)(li + mj) \\ &= 1 - (-l^2 - m^2 + li \cdot mj + mj \cdot li) \\ &= 1 + l^2 + m^2 - lm(i \cdot j - j \cdot i) \\ &= 1 + l^2 + m^2. \end{aligned}$$

Then, by Lemma 6.8, if  $p \mid 1 + l^2 + m^2$ , we must have  $p \mid 1 + li + mj$  or  $p \mid 1 - li - mj$ . But then

$$\frac{1}{p} \pm \frac{li + mj}{p}$$

is a Hurwitz Integer, which is impossible, as  $p > 2$ . Therefore,  $p$  is not a Hurwitz Integer.  $\square$

**Theorem 6.11.** *Let  $p$  be a regular odd prime. Then  $p$  can be written as the sum of four squares.*

*Proof.* Since  $p$  is not a Hurwitz prime, it can be factored as

$$(6.1) \quad p = (a + bi + cj + dk)\alpha.$$

Then

$$\bar{p} = p = \bar{\alpha}(a - bi - cj - dk).$$

Multiplying, we obtain

$$p^2 = (a + bi + cj + dk)\alpha\bar{\alpha}(a - bi - cj - dk) = (a^2 + b^2 + c^2 + d^2)||a||^2.$$

Since  $p$  is not a Hurwitz prime, the original factorization in (1) is nontrivial; therefore, we know that both parts of the factorization

$$p^2 = (a^2 + b^2 + c^2 + d^2)||a||^2$$

are equal to  $p$ . Then  $a^2 + b^2 + c^2 + d^2 = p$ , as desired.  $\square$

Finally, we are ready to prove Lagrange's Four-Square Theorem.

**Theorem 6.12** (Lagrange's Four-Square Theorem). *Every nonnegative integer  $n$  can be written as*

$$n = a^2 + b^2 + c^2 + d^2,$$

*where  $a, b, c$ , and  $d$  are nonnegative integers.*

*Proof.* We start with base cases. We have

$$1 = 0^2 + 0^2 + 0^2 + 1^2$$

and

$$2 = 1^2 + 1^2 + 0^2 + 0^2.$$

By Theorem 6.11, any odd regular prime can be expressed in this form as well. Let

$$n = \prod_{i=1}^k p_i^{e_i}.$$

Since each of the  $p_i$  can be written in the desired form, Lemma 6.7 produces the desired result.  $\square$

## REFERENCES

- [CW11] Erin Compaan and Cynthia Wu. On hurwitz and lipschitz quaternions and lagrange's four-square theorem. *SPWM*, 2011.
- [Fue35] Rud Fueter. Über die analytische darstellung der regularen funktionen einer quaternionenvariablen. *Commentarii Mathematici Helvetici*, 1935.
- [Ham66] William Roward Hamilton. *Elements Of Quaternions*. Longmans, Green, 1866.
- [Sud98] Anthony Sudbery. Quaternionic analysis. *Mathematical Proceedings of the Cambridge Philosophical Society*, 85(2), 1998.